

**INVENTORIES OF BIRDS AND MEDIUM- TO LARGE-BODIED
MAMMALS IN FIVE ATLANTIC FOREST REMNANTS IN SOUTHERN
BRAZIL**

*Juliano A. Bogoni*¹, *Luan G. Araujo Goebel*^{1*}, *Talita C. Bogoni*² & *Manoel Santos-Filho*¹



¹ Universidade do Estado de Mato Grosso, Centro de Pesquisa de Limnologia, Biodiversidade e Etnobiologia do Pantanal; Programa de Pós-Graduação em Ciências Ambientais, Laboratório de Mastozoologia, Cidade Universitária, Av. Santos Dumont s/n, CEP: 78200-000, Cáceres, Mato Grosso, Brazil.

² Instituto Federal Catarinense, Campus Concórdia, Rod. SC-283, Km 8, Vila Fragosos, CEP: 89700-000, Concórdia, Santa Catarina, Brazil.

E-mails: bogoni.ja@gmail.com; lg.araujogoebel@gmail.com (*corresponding author); tbogoni@gmail.com; manoel.filho@unemat.br

SUPPLEMENTARY MATERIAL



Figure S1. Some bird species recorded in five subtropical Atlantic Forest fragments located in Ipumirim municipality, southern Brazil. (A) *Theristicus caudatus* (Boddaert, 1783); (B) *Buteo magnirostris* (Gmelin, 1788); (C) *Paroaria coronata* (Miller, 1776); (D) *Trogon surrucura* (Spix, 1824); (E) *Leptotila verreauxi* Bonaparte, 1855; (F) *Penelope obscura* Temminck, 1815.



Figure S2. Some mammal species recorded in five subtropical Atlantic Forest fragments located in Ipumirim municipality, southern Brazil. (A) *Leopardus guttulus* Hensel, 1872; (B) *Herpailurus yagouaroundi* (É. Geoffroy Saint-Hilaire, 1803); (C) *Hydrochoerus hydrochaeris* (Linnaeus, 1766); (D) *Cerdocyon thous* (Linnaeus, 1766); (E) *Nasua nasua* (Linnaeus, 1766); (F) *Philander frenatus* Olfers, 1818; (G) *Dasypus novemcinctus* Linnaeus, 1758; (H) *Cabassous tatouay* Desmarest, 1804; (I) *Eira barbara* (Linnaeus, 1758); (J) *Didelphis aurita* (Wied-Neuwied, 1826); (K) *Sapajus nigritus* Goldfuss, 1809; (L) *Galictis cuja* (Molina, 1782).